

Circumcision Devices



Circumcision Devices

Disposable Circumcision Anastomat Kit

Manufacturer: Weihai Medison Medical Equipment Co.,Ltd

Production License Number: Lu FDA Production License 20130088

The Registered Address: Qujiahe Industrial Park, Yangting Town, Huancui District, Weihai City, Shandong Province, China



Specifications	Product Composition	Product Model (mm)
Adult	inner ring, auxiliary ring, screws, elastic rope	20、22、24、26、28、30、32、34
Child	stemloop with handle, suture string	11、12、13、14、15、16、17、18、18A、18B、19

■ Disposable Circumcision Anastomat Kit

Medical Equipment Registration Number:
Lu Machinery Registration Approval 20162020372

The Main Product: Disposable Circumcision Anastomat


The Attachment: Iodophor Cotton Swab; Self-adhesive Elastic Bandage and Protective Wound Dressing

Quantity: Disposable Circumcision Anastomat 1 pcs

Iodophor Cotton Swab 10 pcs

Self-adhesive Elastic Bandage 1 roll

Protective Wound Dressing 5 pcs



■ Circumcision Devices (Simple Packing)

Medical Equipment Registration Number:
Lu Machinery Registration Approval 20172020238



■ Product Features:



Traditional surgery: Long operation time, large amount of bleeding, slow recovery, large and irregular wound, requiring suture unsightly appearance.

Laser surgery: Although it has a good hemostatic effect, it still needs to ligate important blood vessels, and the recovery is slow.

Our product: The redundant prepuce falling off easily, fast recovery, no scar after surgery, smooth and beautiful appearance after healing.



The whole operation is simple, safe and painless. The operation only lasts 3-5 minutes, with no surgery and suture free. The patients can get rapid recovery and won't affect the work and life after operation.



The circumcision device can deduce more than 60% infection chance of HIV, playing a certain role in preventing the spread of HIV. With advanced technologies and outstanding advantages, our circumcision device can help male patients of all ages to solve their secret sorrow.



The circumcision device promotes penis to erect frequently and constantly expand at night, in which the cell and tissue of penis is activated, resulting in secondary growth.



After circumcision, epidermal keratinization of glans is caused, which helps desensitization, thus enhancing the sexual function.



It can prevent male urethritis, inflammation of glans and canceration, and meanwhile avoid cross-infection and gynecological diseases.

■ Application

Phimosis and redundant prepuce.

■ Applicable departments

Urology, Surgery, Pediatrics

■ Contraindication

Acute urinary tract infection; concealed penis; redundant patient with shore tie; nervous system disease or mental system disease; leukemia and patient with history of haemorrhagic disease; hypertension, cardiac; STD patient; diabetic.